

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 426 048 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
09.06.2004 Bulletin 2004/24

(21) Application number: 04002911.8

(22) Date of filing: 17.09.1996

(51) Int Cl.7: **A61K 31/192**, A61K 31/4418,
A61K 31/203, A61K 31/426,
A61K 31/4439, A61K 31/455,
A61P 3/06

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**

(30) Priority: 18.09.1995 US 3869 P
06.10.1995 US 4897 P
11.01.1996 US 9884 P
24.05.1996 US 18318 P
10.07.1996 US 21839 P

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
96931613.2 / 0 859 608

(71) Applicant: **LIGAND PHARMACEUTICALS, INC.**
San Diego, CA 92121 (US)

(72) Inventors:
• Heyman, Richard A.
Encinitas CA 92024 (US)
• Cesario, Rosemary
San Diego CA 92117 (US)
• Mukherjee, Ranjan
San Diego CA 92127 (US)

(74) Representative: **Viering, Jentschura & Partner**
Steinsdorfstrasse 6
80538 München (DE)

Remarks:


This application was filed on 10/02/2004 as a
divisional application to the application mentioned
under INID code 62.

(54) **Treating hypertriglyceridemia with RXR agonists in combination with a PPARgamma agonist**

(57) This invention relates to methods and compositions for the treatment of cardiovascular disease, hypertriglyceridemia, low levels of lipoprotein lipase, low

levels of HDL cholesterol and for increasing adipocyte differentiation using an RXR agonist in combination with a PPAR γ agonist such as a thiazolidinedione compound.

FIG. 3.



Description**FIELD OF THE INVENTION**

5 **[0001]** This invention relates to methods and pharmaceutical compounds for treating diabetes and related symptoms.

BACKGROUND OF THE INVENTION

10 **[0002]** Non-insulin-dependent diabetes mellitus (NIDDM, type II diabetes) is characterized by abnormalities in insulin secretion and insulin action. NIDDM constitutes 90-95% of the approximately 6 million diagnosed diabetics in the United States. NIDDM is characterized by hyperglycemia, the result of insulin resistance in peripheral tissues (skeletal muscle and adipose tissue), where insulin-stimulated uptake/utilization of glucose is blunted, and in liver, where insulin suppression of glucose output is insufficient. These impairments in insulin action play an important role in the development of elevated fasting blood glucose and glucose intolerance.

15 **[0003]** Diet and exercise are first-line therapy for NIDDM patients. NIDDM patients also take oral hypoglycemic drugs to control blood glucose levels. The most widely used hypoglycemic agents are various formulations of insulin and sulfonylureas. A major drawback with these therapies is the occurrence of potentially life-threatening hypoglycemia due to hyperinsulinemia.

20 **[0004]** The hyperinsulinemia that can occur with these therapies is also associated with an elevated risk of cardiovascular disease, a major killer of diabetics. Therefore, a need exists for antidiabetic drugs which do not increase circulating insulin concentrations.

25 **[0005]** A new class of compounds, thiazolidinediones, have been documented to effect antihyperglycemic activity by increasing insulin action rather than by promoting insulin secretion. Thiazolidinediones ameliorate insulin resistance and normalize plasma glucose and insulin (where elevated) without causing a hypoglycemic state, even at very high doses. The thiazolidinedione insulin sensitizers, e.g., ciglitazone, englitazone, pioglitazone, BRL 49653 (5-[[4-[2-(methyl-2-pyridinylamino)ethoxy]phenyl]methyl]-2,4-thiazolidinedione), and troglitazone, enhance insulin-mediated suppression of hepatic glucose output and insulin-stimulated glucose uptake and utilization by adipose tissue. Thiazolidinediones also change glucose transporter (e.g. Glut 4) expression to contribute to increased insulin responsiveness.

SUMMARY OF THE INVENTION

[0006] Applicant has found that RXR agonists mimic or enhance the antidiabetic effects of thiazolidinedione compounds. RXR agonists activate the transcriptional activity of RXR/PPAR γ heterodimers, increase insulin stimulated glucose uptake, lower the level of triglyceride, suppress the level of insulin, and increase the level of HDL cholesterol. 35 Two RXR agonists have been shown to lower glucose, triglycerides and insulin levels in two established animal models of NIDDM, i.e. the *ob/ob* and *db/db* mice. Therefore, RXR agonists may be used as insulin sensitizers or insulin mimetics in the treatment of NIDDM and related symptoms.

[0007] In addition, the combination of an RXR agonist and a PPAR γ agonist, such as a thiazolidinedione, achieves synergistic activation of the RXR/PPAR γ heterodimers so as to enhance adipogenic and antidiabetic effects of PPAR γ . 40 In *db/db* mice, the combination of an RXR agonist and a PPAR γ agonist was shown to lower the level of glucose more than individual compounds did.

[0008] Therefore, the present invention relates to methods and compositions for treating a host having NIDDM or insulin resistant diabetes by administering to the host a composition containing a pharmaceutically effective amount of an activator of the RXR/PPAR γ heterodimer, including, but not limited to, an RXR agonist. The host may be a human 45 patient or an animal model of human NIDDM. The compositions of this invention are adapted to cure, improve or prevent one or more symptoms of NIDDM in the host. A preferred drug is highly potent and selective with low toxicity. In this regard, those skilled in the art will recognize NIDDM as an example of a metabolic disease that can be treated with the RXR agonist-containing compounds and compositions of the present invention. Other examples of metabolic diseases treatable with the compounds and compositions of the present invention include, but are not limited to, obesity 50 and thyroid hormone abnormalities.

[0009] By "pharmaceutically effective amount" is meant an amount of a pharmaceutical compound or composition having a therapeutically relevant effect on NIDDM. A therapeutically relevant effect relieves to some extent one or more symptoms of NIDDM in a patient or returns to normal either partially or completely one or more physiological or biochemical parameters associated with or causative of NIDDM, e.g. increasing the sensitivity of cellular response to circulating insulin, curing, reducing, or preventing one or more clinical symptoms of NIDDM, including, but not limited to, hyperglycemia, hyperinsulinemia and hypertriglyceridemia. In a preferred embodiment, a pharmaceutically effective amount of a compound or composition means an amount that increases the uptake of glucose by adipose tissue or muscle tissue. In another preferred embodiment, a pharmaceutically effective amount of a compound or composition 55

means an amount that increases the uptake of triglyceride by adipose tissue.

[0010] By "activator of the RXR/PPAR γ heterodimer" is meant a compound or composition which when combined with the RXR/PPAR γ heterodimer increases the transcriptional regulation activity of the heterodimer, as measured by an assay known to one skilled in the art, including, but not limited to, the "co-transfection" or "cis-trans" assays described or disclosed in U.S. Patent Nos. 4,981,784, 5,071,773, 5,298,429, 5,506,102, WO89/05355, WO91/06677, WO92/05447, WO93/11235, WO95/18380, PCT/US93/04399, PCT/US94/03795 and CA 2,034,220, which are incorporated by reference herein. It includes, but is not limited to, compounds that bind RXR, PPAR γ , or both.

[0011] By "RXR agonist" is meant a compound or composition which when combined with RXR homodimers or heterodimers increases the transcriptional regulation activity of RXR, as measured by an assay known to one skilled in the art, including, but not limited to, the "co-transfection" or "cis-trans" assays described or disclosed in U.S. Patent Nos. 4,981,784, 5,071,773, 5,298,429, 5,506,102, WO89/05355, WO91/0667, WO92/05447, WO93/11235, WO95/18380, PCT/US93/04399, PCT/US94/03795 and CA 2,034,220, which are incorporated by reference herein. It includes, but is not limited to, compounds that preferentially activate RXR over RAR (i.e. RXR specific agonists), and compounds that activate both RXR and RAR (i.e. pan agonists). It also includes compounds that activate RXR in a certain cellular context but not others (i.e. partial agonists). Compounds disclosed or described in the following articles, patents and patent applications which have RXR agonist activity are incorporated by reference herein: U.S. patents 5,399,586 and 5,466,861, WO96/05165, PCT/US95/16842, PCT/US95/16695, PCT/US93/10094, WO94/15901, PCT/US92/11214, WO93/11755, PCT/US93/10166, PCT/US93/10204, WO94/15902 PCT/US93/03944, WO93/21146, provisional applications 60,004,897 and 60,009,884, Boehm, et al. *J. Med. Chem.* 38(16):3146-3155, 1994, Boehm, et al. *J. Med. Chem.* 37(18):2930-2941, 1994, Antras et al., *J. Biol. Chem.* 266:1157-1161 (1991), Salazar-Olivo et al., *Biochem. Biophys. Res. Commun.* 204:157-263 (1994) and Safanova, *Mol. Cell. Endocrin.* 104:201-211 (1994). RXR specific agonists include, but are not limited to, LG 100268 (i.e. 2-[1-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthyl)-cyclopropyl]-pyridine-5-carboxylic acid) and LGD 1069 (i.e. 4-[(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthyl)-2-carbonyl]-benzoic acid), and analogs, derivatives and pharmaceutically acceptable salts thereof. The structures and syntheses of LG 100268 and LGD 1069 are disclosed in Boehm, et al. *J. Med. Chem.* 38(16):3146-3155, 1994, incorporated by reference herein. Pan agonists include, but are not limited to, ALRT 1057 (i.e. 9-cis retinoic acid), and analogs, derivatives and pharmaceutically acceptable salts thereof.

[0012] In a preferred embodiment, the pharmaceutical composition also contains a pharmaceutically effective amount of a PPAR γ agonist. Alternatively, a second composition containing a pharmaceutically effective amount of a PPAR γ agonist is administered to the host separately. In a further preferred embodiment, a compound having agonist activity for both RXR and PPAR γ is used.

[0013] By "PPAR γ agonist" is meant a compound or composition which when combined with PPAR γ increases a reaction typical for the receptor, e.g., transcriptional regulation activity, as measured by an assay known to one skilled in the art, including, but not limited to, the "co-transfection" or "cis-trans" assays described or disclosed in U.S. Patent Nos. 4,981,784 and 5,071,773 and Lehmann, et al., *J. Biol. Chem.* 270:12953-12956 (1995), which are incorporated by reference herein. A preferred PPAR γ agonist is a thiazolidinedione compound, including, but not limited to, BRL 49653, troglitazone, pioglitazone, ciglitazone, WAY-120,744, englitazone, AD 5075, darglitazone, and analogs, derivatives and pharmaceutically acceptable salts thereof. Compounds disclosed in Tontonez et al., *Genes & Develop.* 8:1224-1234 (1994), Tontonez et al., *Cell* 79:1147-1156 (1994), Lehmann et al., *J. Biol. Chem.* 270(22):1-4, 1995, Amri et al., *J. Lipid Res.* 32:1449-1456 (1991), Amri et al., *J. Lipid Res.* 32:1457-1463, (1991) and Grimaldi et al., *Proc. Natl. Acad. Sci. USA* 89:10930-10934 (1992) are incorporated by reference herein.

[0014] In a further preferred embodiment, the pharmaceutical composition also contains a pharmaceutically effective amount of insulin, insulin derivative, insulin secretagogue, insulin sensitizer, or insulin mimetic. Alternatively, a composition containing a pharmaceutically effective amount of insulin, insulin derivative, insulin secretagogue, insulin sensitizer, or insulin mimetic is administered to the host separately.

[0015] A composition containing a pharmaceutically effective amount of an active ingredient may be administered orally or systemically to a host. In a preferred embodiment, it is administered orally.

[0016] In another aspect, this invention features a pharmaceutical composition for treating NIDDM containing a pharmaceutically effective amount of an RXR agonist; and a pharmaceutically acceptable carrier adapted for a host having NIDDM. In a preferred embodiment, the pharmaceutical composition also includes a pharmaceutically effective amount of insulin, insulin derivative, insulin secretagogue, insulin sensitizer, insulin mimetic or PPAR γ agonist.

[0017] In a preferred embodiment, the composition is held within a container which includes a label stating to the effect that the composition is approved by the FDA in the United States (or an equivalent regulatory agency in a foreign country) for treating NIDDM or for treating hyperglycemia, hyperinsulinemia or hypertriglyceridemia. Such a container provides a therapeutically effective amount of the active ingredient to be administered to a host.

[0018] In another aspect, this invention features methods for screening for candidate compounds useful for treating NIDDM. These methods select compounds or compositions which when combined with the RXR/PPAR γ heterodimer increase the transcriptional regulation activity of the heterodimer, as measured by an assay known to one skilled in

the art, including, but not limited to, the "co-transfection" or "cis-trans" assays described or disclosed in U.S. Patent Nos. 4,981,784, 5,071,773, 5,298,429, 5,506,102, WO89/05355, WO91/06677, WO92/05447, WO93/11235, WO95/18380, PCT/US93/04399, PCT/US94/03795 and CA 2,034,220, which are incorporated by reference herein. In one example, a candidate compound such as a potential RXR agonist is administered to an adipocyte or a preadipocyte. The level of lipid in the cell is measured, and an increased accumulation of lipid after the treatment with the candidate compound indicates that the candidate compound is useful for treating NIDDM. In preferred embodiments, the level of lipid is measured by oil red O staining or detecting the level of triglyceride in the cell.

[0019] In another example, a candidate compound such as a potential RXR agonist is administered to an adipocyte or a preadipocyte and the transcription level of an adipocyte specific gene (e.g. lipoprotein lipase gene or PPAR γ gene) is measured. An increased transcription of the adipocyte specific gene after the treatment with the candidate compound indicates that the candidate compound is useful for treating NIDDM.

[0020] In yet another example, a candidate compound such as a potential RXR agonist is administered to an adipocyte or a preadipocyte and the level of glucose uptake is measured. An increased glucose uptake after the treatment with the candidate compound indicates that the candidate compound is useful for treating NIDDM. Alternatively, both the candidate compound and insulin are administered to the cell and the level of glucose uptake is compared to that in the same cell treated with insulin alone. A higher level of glucose uptake in the cell treated by the candidate compound and insulin indicates that the candidate compound is an insulin sensitizer and useful for treating NIDDM.

[0021] Other features and advantages of the invention will be apparent from the following detailed description of the invention, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022]

Figure 1a is a graph showing the extent of adipocyte differentiation in 3T3-L1 cells as measured by the levels of triglyceride in 3T3-L1 preadipocytes treated with various combinations of an RXR agonist (LG 100268), a PPAR γ agonist (thiazolidinedione compound BRL 49653), and insulin. The retinoid and BRL 49653 were used at 1 μ M.

Figure 1b is a graph showing the extent of adipocyte differentiation in 3T3-L1 cells as measured by the level of triglyceride in 3T3-L1 preadipocytes treated with various combinations of an RXR agonist (LGD 1069), a PPAR γ agonist (thiazolidinedione compound BRL 49653), and insulin. The retinoid and BRL 49653 were used at 1 μ M.

Figure 2a is a graph showing the level of LPL mRNA in 3T3-L1 cells treated with various combinations of an RXR agonist (LG 100268), a PPAR γ agonist (thiazolidinedione compound BRL 49653), and insulin. The retinoid and BRL 49653 were used at 1 μ M.

Figure 2b is a graph showing the level of PPAR γ mRNA in 3T3-L1 cells treated with various combinations of an RXR agonist (LG 100268), a PPAR γ agonist (thiazolidinedione compound BRL 49653), and insulin. The retinoid and BRL 49653 were used at 1 μ M.

Figure 3 is a graph showing the level of glucose in *db/db* mice treated with LG 100268, BRL 49653, and a combination of LG 100268 and BRL 49653, respectively.

Figure 4 is a graph showing the level of triglycerides in *db/db* mice treated with LG 100268, BRL 49653, and a combination of LG 100268 and BRL 49653, respectively.

Figure 5 is a graph showing the level of HDL cholesterol in *db/db* mice treated with LG 100268, BRL 49653, and a combination of LG 100268 and BRL 49653, respectively.

Figure 6 is a graph showing the level of triglycerides in *ob/ob* mice treated with LGD 1069, LG 100268, and BRL 49653, respectively.

Figure 7 is a graph showing the level of glucose in *ob/ob* mice treated with LGD 1069, LG 100268, and BRL 49653, respectively.

Figure 8 is a graph showing the level of insulin in *ob/ob* mice treated with LGD 1069, LG 100268, and BRL 49653, respectively.

DETAILED DESCRIPTION OF THE INVENTION

TZD achieves antidiabetic and adipogenic effects through PPAR γ

[0023] Thiazolidinediones are insulin sensitizers that significantly reduce glucose and lipid levels in animal models of NIDDM and obesity (Kees, et al., *J. Medicinal Chem.* 38(4):617-628, 1995; Willson, et al., *J. Medicinal Chem.* 39(3):665-668, 1996; Young, et al. *Diabetes* 44:1087-1092, 1995). Thiazolidinediones improve glucose utilization without stimulating insulin release.

[0024] For example, repeated administration of BRL 49653 to obese mice improves glycemic control by increasing

insulin responsiveness of target tissues. BRL 49653 potentiates insulin-stimulated glucose transport in adipocytes from insulin-resistant obese mice, both by increasing insulin receptor number and by facilitating translocation of GLUT4, from an expanded intracellular pool, to the cell surface.

[0025] Thiazolidinediones are also selective PPAR γ agonists (Lehmann, et al. *J. Biol. Chem.* 270(22):1-4, 1995). Comparison of the EC₅₀ for activation of PPAR γ with the minimum effective dose (MED) for anti-hyperglycemic activity revealed a significant correlation. The correlation between *in vitro* PPAR γ activity and *in vivo* antihyperglycemic activity of thiazolidinediones implicates PPAR γ as the molecular target for the antidiabetic effects of thiazolidinediones.

[0026] PPAR γ is a member of the nuclear receptor superfamily of ligand-activated transcription factors. It is expressed in an adipose-specific manner and its expression is induced early during the differentiation of several preadipocyte cell lines. Forced expression of PPAR γ in fibroblasts resulted in adipocyte differentiation.

[0027] In addition to insulin sensitizing activity, thiazolidinediones have marked adipogenic effects on preadipocyte and mesenchymal stem cells (Tontonoz, et al., *Cell* 79:1147-1156, 1994). Treatment of C3H/10T1/2 cells with BRL 49653 resulted in efficient adipocyte differentiation, showing that ligand-mediated activation of PPAR γ is sufficient to initiate the adipogenic signaling cascade in a mesenchymal stem cell line. PPAR γ is the molecular target for the adipogenic effects of thiazolidinediones.

[0028] Adipogenesis plays a role in the development of NIDDM, which is characterized by not only unbalanced glucose homeostasis, but also elevated levels of circulating lipids. Increases in lipid levels have been shown to interfere with glucose disposal.

[0029] Adipocytes are highly specialized cells that play a critical role in lipid metabolism and energy homeostasis. Their primary role is to store triglycerides in times of caloric excess and to mobilize this reserve during periods of nutritional deprivation.

[0030] Adipocyte differentiation is characterized by a coordinate increase in adipocyte-specific gene expression. PPAR γ is specifically expressed in adipocytes. Its expression is induced early during the course of differentiation of several preadipocyte cell lines. Forced expression of PPAR γ in fibroblasts resulted in adipocyte differentiation.

Synergistic effects of RXR and PPAR γ on adipogenesis

[0031] Expression of PPAR γ is induced early during the differentiation of cultured adipocyte cell lines and is expressed at very high levels specifically in adipose tissue. PPAR γ regulates adipogenesis by modulating the transcription of other adipocyte-specific genes, e.g. adipocyte P2 gene (aP2 gene). aP2 gene encodes an intracellular lipid-binding protein and is expressed exclusively in adipose cells.

[0032] A 518-bp DNA fragment from the 5'-flanking region of the aP2 gene has been identified as an enhancer that directs high-level adipocyte-specific gene expression in both cultured cells and transgenic mice. A pair of elements in the aP2 enhancer, ARE6 and ARE7, bind a nuclear factor termed ARF6 that is detected only in nuclear extracts derived from adipocytes. The ARF6-binding sites are both necessary and sufficient for adipocyte-specific expression, suggesting that the trans-acting factor ARF6 functions as a differentiation-dependent and tissue-specific switch for the aP2 enhancer (Tontonoz, et al. *Genes & Development*, 8:1224-1234, 1994).

[0033] The ARF6 recognition sequence resembles a type of nuclear hormone receptor-binding site known as DR-1 (direct repeat with 1-nucleotide spacer). This motif has been shown to preferentially bind heterodimers of RXR and COUP-TF and heterodimers of RXR and the PPARs. DNA mobility retardation experiments using various HRE sequences as competitor demonstrated that ARF6 preferentially recognizes DR-1 sites.

[0034] ARF6 has been identified as a heterodimeric complex of RXR α and PPAR γ . It has been shown that PPAR γ and RXR α form heterodimers on ARF6-binding sites *in vitro*. Forced expression of these factors in transient transfections is sufficient to activate the adipocyte-specific aP2 enhancer in nonadipose cells such as fibroblasts. This activation is potentiated by peroxisome proliferators, fatty acids, and 9-cis retinoic acid. Antiserum to RXR α specifically inhibits ARF6 activity in adipocyte nuclear extracts.

[0035] Cotransfection of the RXR α expression vector and the PPAR γ expression vector has a synergistic effect to activate the aP2 enhancer in nonadipose cells. Maximal activation of the aP2 enhancer is observed when both PPAR γ , RXR α and their agonists are present.

[0036] Without being bound by any theory, applicant proposes that an RXR agonist affects glucose usage in tissues through the synergistic effects of RXR and PPAR γ heterodimers. The RXR/PPAR γ heterodimers, when activated by an RXR agonist or a combination of an RXR agonist and a PPAR γ agonist, induce adipogenesis and modulates the levels of glucose and triglyceride uptake. Alternatively or in addition, the RXR/PPAR γ heterodimers, when activated by an RXR agonist or a combination of an RXR agonist and a PPAR γ agonist, regulate signaling molecules secreted by adipose tissue such as tumor necrosis factor- α or leptin, which in turn modulates glucose metabolism in other tissues.

Using RXR agonists to mimic or enhance the antidiabetic effects of thiazolidinediones

[0037] PPAR α , β and γ all form heterodimers with RXRs. These RXR/PPAR heterodimers bind to DNA and regulate transcription activity. RXR activators cooperate with PPAR α activators to activate the activity of PPAR α protein (Kliwer, *et al.* Nature 358:771-774 (1992) and Mukherjee, *et al.* Steroid Biochem. Molec. Biol. 51:157-166 (1994)).

[0038] In this invention, a similar synergistic activation was observed with a PPAR γ activator and several RXR activators.

[0039] According to this invention, RXR agonists, e.g. LGD 1069, ALRT 1057 and LG 100268, may be utilized in the treatment of diabetes. We examined four independent parameters for the effects of RXR agonists, i.e. morphological changes, lipid accumulation, regulation of gene expression and increased glucose uptake.

[0040] Two pre-adipocyte cell lines were used to test the theory of RXR activation in the PPAR γ /RXR heterodimer. 3T3-L1 and C3H/10T1/2 cells were obtained from ATCC and are derived from mouse embryo. They are contact inhibited and can be induced to differentiate into adipocyte cells containing large lipid droplets within the cytoplasm. Adipocyte differentiation can be observed by oil red O staining which stains the lipid droplets within the cytoplasm red. The extent of adipocyte differentiation can then be monitored by microscope observation.

[0041] For a more quantitative assay and rapid screening of compounds, a 96-well plate assay was developed to quantitate the amount of triglyceride produced by the differentiating adipocytes. In this assay, cells are grown as a monolayer to confluence on a 96-well plate and treated with BRL 49653, insulin, and retinoids alone or in various combinations.

These treatments induce differentiation to different extent in both 3T3-L1 and C3H/10T1/2 cells. The level of triglyceride accumulation can then be measured via an enzymatic color reaction which can be read in a plate reader.

[0042] A third measure of adipocyte differentiation is to examine regulation of gene expression. The mRNA expression levels of both PPAR γ and lipoprotein lipase (LPL) have been shown to be modulated during adipocyte differentiation. Northern blot analysis was used to dissect the molecular aspects of how retinoids effect target genes of differentiating adipocytes. PPAR γ , lipoprotein lipase (LPL), and β -actin (loading control) mRNA levels were monitored after cells were treated with thiazolidinediones and retinoids.

[0043] A fourth indicator for the utility of a compound in treating NIDDM or insulin-resistant diabetes is the compound's ability to enhance insulin-stimulated glucose uptake. Labeled 2-deoxyglucose (2-DOG, a glucose analog) assay was performed with a preadipocyte cell line in the presence of insulin and a candidate compound to measure the level of 2-DOG incorporation.

A) Retinoid Modulation of Lipid Accumulation in 3T3-L1 Cells Stained with Oil Red O

[0044] Table 1 shows the percent of 3T3-L1 cells that had differentiated into adipocytes as observed by oil red O staining assay. BRL 49653 and LG 100268 were used at 1 μ M, insulin was used at 0.01 mg/ml. Wells treated with LG 100268 had bigger redder lipid droplets within the cytoplasm.

[0045] BRL 49653 and LG 100268 treatment alone induced 50% of the cells to differentiate into adipocytes. This was increased dramatically with the addition of insulin. Insulin in combination with BRL induced 80% of the 3T3-L1 cells into adipocytes while the combination of insulin and LG 100268 induced 90% of the cells to differentiate. When BRL 49653 was used in combination with LG 100268, an RXR agonist, the amount of adipocytes differentiation was also increased dramatically. Other RXR agonists mimic the activity of LG 100268. For example, the addition of ALRT 1057 (pan agonist) or LGD 1069 (RXR specific agonist) in combination with BRL 49653 increased the amount of differentiation, albeit to a lesser extent than the strong RXR agonist LG 100268. The combination of insulin with BRL 49653 and LGD 1069 had a strong differentiating effect (95%) on the 3T3-L1 cells.

B) Retinoid Modulation of Triglyceride Content in Differentiated Adipocytes

[0046] The retinoid modulation of lipid formation was quantitated by monitoring triglyceride formation. Figures 1a and 1b show triglyceride accumulation in 3T3-L1 cells treated with a retinoid (LG 100268 or LGD 1069) alone or in combination with a thiazolidinedione and insulin. Retinoids and BRL 49653 were used at 1 μ M, insulin was used at 0.01 mg/ml for all experimental combinations.

[0047] Insulin, BRL 49653 and retinoids all induced some triglyceride accumulation when used alone, with LG 100268 giving the largest response. The addition of retinoids (LGD 1069, LG 100268) with the thiazolidinedione (BRL 49653) to the assay increased the amount of triglyceride accumulation in 3T3-L1 differentiating adipocytes. This was also observed when BRL 49653 or LG 100268 was used in combination with insulin. The largest accumulation of triglyceride was seen in the cells treated with LG 100268, BRL 49653 and insulin together. Similar results were observed when LGD 1069 replaced LG 100268 in the study. These results concur with those obtained in the oil red O staining assays.

C) PPAR γ and LPL mRNA modulation in differentiating 3T3-L1 cells

[0048] Adipocyte specific genes were monitored via northern blot analysis. Figure 2a and 2b show the expression pattern of LPL (lipoprotein lipase) mRNA and PPAR γ mRNA in cells that were treated for 7 days with BRL 49653 (1 μ M), LG 100268 (1 μ M) and insulin (0.01 mg/ml), alone or in combination.

[0049] Northern blot analysis shows an increase in the relative signal normalized to β -actin of both LPL and PPAR γ mRNA expression in cells treated with either compound alone. There was a three to five fold increase in mRNA levels for these adipocyte target genes demonstrating that transcriptional regulation occurs with treatment by insulin, BRL 49653 and LG 100268. Combination of insulin, BRL 49653 and LG 100268 did not further enhance the mRNA level.

[0050] These data demonstrate that, in 3T3-L1 cells, RXR agonists induce adipocyte differentiation by themselves or in combination with thiazolidinediones or insulin. RXR agonists enhance the activity of thiazolidinediones and insulin. Three independent measurements support that RXR agonists contribute to the modulation of the RXR/PPAR γ heterodimer in regulating adipocyte differentiation and useful in treating NIDDM.

D) LG 100268 enhances insulin stimulated glucose uptake in 3T3-L1 cells

[0051] A murine preadipocyte cell line, 3T3-L1, is widely used to study glucose uptake, adipogenesis, and has been used in the characterization of thiazolidinediones and other PPAR γ activators. Insulin stimulated uptake of labeled 2-deoxyglucose (2-DOG, a glucose analog) was observed in 3T3-L1 cells treated with BRL 49653 or LG 100268.

[0052] 3T3-L1 cells were treated with BRL 49653 (10 μ M) or LG 100268 (1 μ M) for 10 days. Insulin was added to the cells at a concentration of 0.01 mg/ml for 5 days, thereafter no insulin was added. Labeled 2-deoxyglucose uptake assay was performed (Szalkowski, et al., J. Endocrin. 136:1474-1481, 1995). The level of incorporated labeled 2-DOG was normalized to the amount of total cellular protein. About 5-fold increase in 2-DOG uptake was observed in 3T3-L1 cells treated with BRL 49653 alone. About 2-fold increase in 2-DOG uptake was observed in 3T3-L1 cells treated with LG 100268 alone.

[0053] This experiment shows that an RXR agonist increases insulin mediated glucose uptake in 3T3-L1 cells like a known insulin sensitizer, a thiazolidinedione. It demonstrates directly that RXR agonists can be used to treat a major symptom of NIDDM, i.e. insulin resistance. Other RXR agonists useful for the treatment of NIDDM can be confirmed using this assay.

E) Retinoid Modulation of Lipid Accumulation in C3H/10T1/2 Cells Stained with Oil Red O

[0054] To further assess the retinoid regulation of adipocyte function we examined C3H/10T1/2 cells, which are mouse embryo fibroblast/multipotent stem cells that can be induced to differentiate into adipocytes or muscle cells. Table 2 shows the percent of C3H/10T1/2 cells that have differentiated into adipocytes as observed by oil red O staining assay. Experiment A was conducted in the presence of BRL 49653. Experiment B was conducted in the presence of both BRL 49653 and insulin. ALRT 1057 and LGD 1069 were used at 1 μ M and TTNPB (RAR selective compound) was used at 10 nM concentration. Insulin was used at 0.01 mg/ml; and BRL 49653 was used at 0.1, 1 and 10 μ M. C3H/10T1/2 cells were treated for 7 days, cells were stained with oil red O and observed under the microscope.

[0055] Morphological changes were observed when the cells were treated with retinoids alone, even though the cells did not fully differentiate into adipocytes. BRL 49653 alone caused no more than 10 percent of the cells to undergo adipocyte differentiation. However, when BRL 49653 was used in combination with an RXR agonist, LGD 1069 or ALRT 1057, there was a large increase in the amount of differentiation, the same effect was seen when BRL 49653 was used in combination with insulin.

[0056] The RAR agonist, TTNPB (which does not activate RXR), did not have the same effect as RXR agonist LGD 1069 and ALRT 1057. In fact, TTNPB inhibited the differentiation induced by BRL 49653 or BRL/insulin.

[0057] The above experiments show that a thiazolidinedione (BRL 49653) alone induces a minimal amount of differentiation in C3H/10T1/2 cells. However, when BRL 49653 is used in combination with a retinoid such as LG 100268, LGD 1069, or ALRT 1057, which are RXR agonists, differentiation is dramatically increased. This is not seen with the pure RAR agonist, TTNPB. These data support that the PPAR γ /RXR heterodimers, which drive the adipocyte differentiation process, can be activated and enhanced by the binding of an RXR agonist. RXR agonists are useful for modulating the levels of glucose and triglyceride uptake.

Using RXR agonists to lower levels of glucose and triglycerides in animal models of NIDDM(A) In vivo experiment with *db/db* mice

5 [0058]

Animals: Strain Diabetic C57BLKS/J - *m +/-db*, 82 mice
 Source: Jackson Lab
 Stock number 000642
 10 Genotype *m +/-db xm +/-db*
 DOB 6/19/96±3d, DOA 7/23/96 - 34d old
 Date of study: 8/5/96 - 8/21/96, 44-63d old

15 [0059] Mice were identified by an ear punch code (#1-82) and separated into 8 groups (A-H). Each group consisted of 10 mice separated into 4 cages with 2-3 mice/cage. Several mice were lost during the course of the study from fluid injections into the lungs resulting in 2 groups with 9 mice/group. Control group C consisted of 12 mice. The mice were fed pelleted Purina Lab chow #5015 containing 3.83kcal/g with a caloric composition of 56% carbohydrate, 26% fat and 18% protein. Food and water were provided ad libitum. Food intake/cage was measured over selected periods and expressed as g food consumed/100g mouse/day.

20 [0060] On days of study, food was removed from the cages at selected intervals of time between 6:15 and 7:00 AM. Animal weights were recorded 2h after start of fast and blood samples taken after a 3h fast. Blood was drawn from a cut at the tip of the tail and collected into a heparinized capillary tube (approx. 75µl volume). After centrifugation, the hematocrit was read in a microcapillary reader, recorded and the tube broken for recovery of plasma for analysis of glucose, triglyceride and insulin concentration.

25 [0061] Blood samples were collected on days 0, 1, 3, 7, 10, and on the final day of study, days 13-15. After collection of blood samples on day 0, animals were refed their Purina chow diet and subsequently gavaged with control solution in group C and one of seven test solutions in groups designated A, B, D-H. The volume administered was equivalent in each group averaging .6ml/42g mouse (.01429 ml/g). The various solutions were gavaged daily to their respective groups based on the animals weight taken that morning or the weight of the animal taken on the preceding day of weighing. To assess alterations in plasma FFA, on day 10 an additional 75 µl blood sample was collected into an EDTA-coated capillary tube immediately after collecting the basal heparinized capillary tube sample.

30 [0062] On the final day of study, mice were not gavaged with test solution. The last gavage was administered on the day preceding the final day, *i.e.*, day 12 for animals terminated on day 13, day 13 for animals terminated on day 14 and day 14 for animals terminated on day 15. Terminal blood was collected by decapitation to provide serum for assessment of HDL.

Results:

40 [0063] Figure 3 shows that BRL49653 and LG100268 each independently lowered the level of glucose in *db/db* mice. In addition, the combination of BRL49653 and LG100268 lowered the glucose level more than each compound did by itself. BRL 49653 (1 mg/kg) and LG 100268 (20 mg/kg) lowered the level of glucose by 40% by day 15 in comparison to control. The thiazolidinedione BRL 49653 at 1 mg/kg showed similar efficacy. The combination of an RXR activator and a PPAR activator showed greater efficacy, leading to almost 50% drop in the level of glucose. The effect of the combination was rapid, glucose levels were reduced by day 1 of the study and reached a steady state by day 4. This indicates a rapid resetting of the steady state levels of glucose homeostasis. Therefore, RXR activators enhance the efficacy of PPARγ activators, and vice versa.

45 [0064] Figure 4 shows that RXR activators lowered the level of triglycerides in *db/db* mice. LG 100268 (20 mg/kg) lowered the level of triglycerides 40% by day 15 of the study. BRL 49653 (1 mg/kg) showed similar efficacy. The combination of these two compounds worked even better.

50 [0065] Figure 5 shows that RXR activators modulators increased the level of HDL cholesterol in *db/db* mice. LG 100268 (20 mg/kg) increased HDL-cholesterol levels (20%) in comparison to controls in *db/db* mice. BRL 49653 caused an equivalent level of increase. The combination of these two compounds showed a higher increase. HDL-C was measured by the precipitation method using kits obtained from Bohringer-Mannheim (catalog # 543004 and 427578).

55

(B) In vivo experiment with *ob/ob* mice**[0066]**

Animals: Strain Obese C57 BL/6J-*Lep^{ob(4)}*, 121 mice
 Source: Jackson Lab Stock number 000632
 Genotype *Lep^{ob(4)}/+* x *Lep^{ob(4)}/+*
 DOB 5/22/96±3d, DOA 7/2/96 - 41d old
 Date of study: 7/21/96 - 8/2/96 - 49-70d old

[0067] Mice were identified by an ear punch code (#1-121) and separated into 12 groups. Each group consisting of 10 mice. (one mouse was terminated before start of the study because of bad teeth causing initial weight loss. As in preceding studies, mice in each group were housed in 4 cages (2-3 mice/cage) and provided water and Purina Lab chow #50515 *ad libitum*.

[0068] On days of study, food was removed from the cages at selected intervals of time between 6:15 and 7:00 AM. Animal weights were recorded 2h after start of fast and blood samples taken after a 3h fast. Blood was drawn from a cut at the tip of the tail and collected into a heparinized capillary tube (approx. 75µl volume). After centrifugation, the hematocrit was read in a microcapillary reader, recorded and the tube broken for recovery of plasma for analysis of glucose, triglyceride and insulin concentration.

[0069] Blood samples were collected on days 0, 3, 6, 8, 10, 14±1 and final collections were made on days 15, 16 or 17. FFA samples were collected on day 10 by drawing a second 75µl blood sample in a non-heparinized tube coated with EDTA. This tube was collected immediately after the heparinized tube collection.

[0070] Animals in groups H were administered control gavage solution (0.6 ml/42 g) daily commencing on day 0 and ending on the day preceding the final day. 11 test solutions were administered to animals in groups designated A-G, I-L. All gavage solutions were administered after refeeding on days in which mice were fasted for blood sampling.

Results:

[0071] Figure 6 shows that RXR modulators lowered the level of triglycerides in *ob/ob* mice. RXR activators LGD 1069 (30 mg/kg) and LG 100268 (20 mg/kg) lowered the level of triglycerides by 34% and 60% respectively in *ob/ob* mice by day 14 of the study. BRL 49653 was also able to lower the level of triglycerides, although not as efficacious as LG100268.

[0072] Figure 7 shows that RXR modulators lowered the level of glucose in *ob/ob* mice. RXR activators LGD 1069 at 30 mg/kg and LG 100268 at 20 mg/kg lowered the level of glucose nearly 50% in comparison to control. The level of glucose was reduced to almost euglycemic levels by day 14 of the study. BRL 49653 (0.4 mg/kg) showed similar efficacy.

[0073] Figure 8 shows that RXR modulators, LGD 1069 (30 mg/kg) and LG 100268 (20 mg/kg), lowered the level of insulin in *ob/ob* mice. LG100268 lowered the level of insulin by 66% by day 14 of the study. LGD 1069 showed lower efficacy.

There was a very rapid effect of the compounds since the level of insulin started to drop in day 1.

Pharmaceutical Formulations and Modes of Administration

[0074] The particular compound that affects the disorders or conditions of interest can be administered to a patient either by themselves, or in pharmaceutical compositions where it is mixed with suitable carriers or excipient(s). In treating a patient exhibiting a disorder of interest, a therapeutically effective amount of a agent or agents such as these is administered. A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or a prolongation of survival in a patient.

[0075] The compounds also can be prepared as pharmaceutically acceptable salts. Examples of pharmaceutically acceptable salts include acid addition salts such as those containing hydrochloride, sulfate, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, cyclohexylsulfamate and quinate. (See *e.g.*, PCT/US92/03736). Such salts can be derived using acids such as hydrochloric acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, *p*-toluenesulfonic acid, cyclohexylsulfamic acid, and quinic acid. These salts can be prepared by standard techniques. For example, the free base form of the compound is first dissolved in a suitable solvent such as an aqueous or aqueous-alcohol solution, containing the appropriate acid. The salt is then isolated by evaporating the solution. In another example, the salt is prepared by reacting the free base and acid in an organic solvent.

[0076] Carriers or excipients can be used to facilitate administration of the compound, for example, to increase the solubility of the compound. Examples of carriers and excipients include calcium carbonate, calcium phosphate, various sugars or types of starch, cellulose derivatives, gelatin, vegetable oils, polyethylene glycols and physiologically compatible solvents.

5 [0077] In addition, the molecules tested can be used to determine the structural features that enable them to act on the RXR/PPAR γ heterodimer, and thus to select molecules useful in this invention. Those skilled in the art will know how to design drugs from lead molecules, using techniques such as those disclosed in PCT publication WO 94/18959, incorporated by reference herein.

10 [0078] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compounds which exhibit large therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. Levels in plasma may be measured, for example, by HPLC.

15 [0079] The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g. Fingl et al., in The Pharmacological Basis of Therapeutics, 1975, Ch. 1 p. 1). It should be noted that the attending physician would know how to and when to terminate, interrupt, or adjust administration due to toxicity, or to organ dysfunctions. Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate (precluding toxicity). The magnitude of an administered dose in the management of the disorder of interest will vary with the severity of the condition to be treated and to the route of administration. The severity of the condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the dose and perhaps dose frequency, will also vary according to the age, body weight, and response of the individual patient. A program comparable to that discussed above may be used in veterinary medicine.

20 [0080] Depending on the specific conditions being treated, such agents may be formulated and administered systemically or locally. Techniques for formulation and administration may be found in Remington's Pharmaceutical Sciences, 18th ed., Mack Publishing Co., Easton, PA (1990). Suitable routes may include oral, rectal, transdermal, vaginal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections, just to name a few.

25 [0081] For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For such transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

30 [0082] Use of pharmaceutically acceptable carriers to formulate the compounds herein disclosed for the practice of the invention into dosages suitable for systemic administration is within the scope of the invention. With proper choice of carrier and suitable manufacturing practice, the compositions of the present invention, in particular, those formulated as solutions, may be administered parenterally, such as by intravenous injection. The compounds can be formulated readily using pharmaceutically acceptable carriers well known in the art into dosages suitable for oral administration. Such carriers enable the compounds of the invention to be formulated as tablets, pills, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

35 [0083] Agents intended to be administered intracellularly may be administered using techniques well known to those of ordinary skill in the art. For example, such agents may be encapsulated into liposomes, then administered as described above. Liposomes are spherical lipid bilayers with aqueous interiors. All molecules present in an aqueous solution at the time of liposome formation are incorporated into the aqueous interior. The liposomal contents are both protected from the external microenvironment and, because liposomes fuse with cell membranes, are efficiently delivered into the cell cytoplasm. Additionally, due to their hydrophobicity, small organic molecules may be directly administered intracellularly.

40 [0084] Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. The preparations formulated for oral administration may be in the form of tablets, dragees, capsules, or solutions. The pharmaceutical compositions of the present invention may be manufactured in a

manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levitating, emulsifying, encapsulating, entrapping or lyophilizing processes.

[0085] Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

[0086] Pharmaceutical preparations for oral use can be obtained by combining the active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

[0087] Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

[0088] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. Liposomes may be used for encapsulated delivery.

[0089] Pharmaceutical formulations disclosed or described in Boehm, et al., WO94/15902 are incorporated by reference herein.

[0090] All publications referenced are incorporated by reference herein, including the nucleic acid sequences and amino acid sequences listed in each publication. All the compounds disclosed and referred to in the publications mentioned above are incorporated by reference herein, including those compounds disclosed and referred to in articles cited by the publications mentioned above.

[0091] Other embodiments of this invention are disclosed in the following claims.

TABLE 1:

Oil Red O Staining in 3T3-L1 Differentiating Adipocytes	
Treatment	Percent of Differentiated Cells
Control	0.1
BRL 49653	50
Insulin	20
LG 100268	50
BRL+Insulin	80
BRL+LG 100268	90
LG 100268+Insulin	90

TABLE 2:

Retinoid Modulation of Adipocyte differentiation in C3H/10T1/2 Cells		
Treatment	Experiment A (+BRL) % differentiated cells	Experiment B (+BRL+Insulin) % differentiated cells
BRL (10 μ M)	10	80
BRL (1 μ M)	8	80
BRL (0.1 μ M)	5	60

TABLE 2: (continued)

Retinoid Modulation of Adipocyte differentiation in C3H/10T1/2 Cells		
Treatment	Experiment A (+BRL) % differentiated cells	Experiment B (+BRL+Insulin) % differentiated cells
BRL (10 μ M)	10	80
+ALRT 1057 (1 μ M)	80	99
+LGD 1069 (1 μ M)	80	99
+TTNPB(10nM)	5	50
BRL (1 μ M)	8	80
+ALRT 1057 (1 μ M)	65	90
+LGD 1069 (1 μ M)	65	90
+TTNPB(10nM)	2	30
BRL (0.1 μ M)	5	60
+ALRT 1057 (1 μ M)	45	85
+LGD 1069(1 μ M)	45	90
+TTNPB (10nM)	0.2	
Undifferentiated controls 0% differentiation.		

Claims

1. Use of a Retinoid X Receptor (RXR) agonist in the manufacture of a pharmaceutical composition for treating a host having cardiovascular disease or preventing cardiovascular disease in a host, wherein said composition further comprises a peroxisome proliferator-activated receptor γ (PPAR γ) agonist, said PPAR γ agonist being a thiazolidinedione compound, wherein said RXR agonist is 2-[1-3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthyl)-cyclopropyl]-pyridine-5-carboxylic acid (LG 100268) or 4-[1,5,6,7,8-tetrahydro-3,5,5,8,8-pentamethyl-2-naphthalenyl) ethenyl] benzoic acid (LGD 1069).
2. Use of a Retinoid X Receptor (RXR) agonist in the manufacture of a pharmaceutical composition for treating a host having hypertriglyceridemia or preventing hypertriglyceridemia in a host, wherein said composition further comprises a peroxisome proliferator-activated receptor γ (PPAR γ) agonist, said PPAR γ agonist being a thiazolidinedione compound, wherein said RXR agonist is 2-[1-3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthyl)-cyclopropyl]-pyridine-5-carboxylic acid (LG 100268) or 4-[1,5,6,7,8-tetrahydro-3,5,5,8,8-pentamethyl-2-naphthalenyl) ethenyl] benzoic acid (LGD 1069).
3. Use of a Retinoid X Receptor (RXR) agonist in the manufacture of a pharmaceutical composition for treating a host having low levels of lipoprotein lipase, wherein said composition further comprises a peroxisome proliferator-activated receptor γ (PPAR γ) agonist, said PPAR γ agonist being a thiazolidinedione compound, wherein said RXR agonist is 2-[1-3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthyl)-cyclopropyl]-pyridine-5-carboxylic acid (LG 100268) or 4-[1,5,6,7,8-tetrahydro-3,5,5,8,8-pentamethyl-2-naphthalenyl) ethenyl] benzoic acid (LGD 1069).
4. Use of a Retinoid X Receptor (RXR) agonist in the manufacture of a pharmaceutical composition for treating a host having low levels of HDL cholesterol, wherein said composition further comprises a peroxisome proliferator-activated receptor γ (PPAR γ) agonist, said PPAR γ agonist being a thiazolidinedione compound, wherein said RXR agonist is 2-[1-3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthyl)-cyclopropyl]-pyridine-5-carboxylic acid (LG 100268) or 4-[1,5,6,7,8-tetrahydro-3,5,5,8,8-pentamethyl-2-naphthalenyl) ethenyl] benzoic acid (LGD 1069).
5. Use of a Retinoid X Receptor (RXR) agonist in the manufacture of a pharmaceutical composition for increasing adipocyte differentiation in a host, wherein said composition further comprises a peroxisome proliferator-activated receptor γ (PPAR γ) agonist, said PPAR γ agonist being a thiazolidinedione compound, wherein said RXR agonist is 2-[1-3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydro-2-naphthyl)-cyclopropyl]-pyridine-5-carboxylic acid (LG 100268) or 4-[1,5,6,7,8-tetrahydro-3,5,5,8,8-pentamethyl-2-naphthalenyl) ethenyl] benzoic acid (LGD 1069) for increasing

adipocyte differentiation

6. The use according to any of claims 1 to 5, wherein said thiazolidinedione compound is selected from the group consisting of BRL 49653, troglitazone, pioglitazone, ciglitazone, WAY-120,744, englitazone, AD 5075 and darglitazone.

7. The use according to claim 6, wherein said composition further comprises insulin, insulin derivative, insulin secretagogue, insulin sensitizer or insulin mimetic.

8. A pharmaceutical composition for the treatment or prevention of cardiovascular disease comprising:

- (a) an RXR agonist;
- (b) a PPAR γ agonist; and
- (c) a pharmaceutically acceptable carrier,

wherein said PPAR γ agonist is a thiazolidinedione compound, and wherein said RXR agonist is selected from the group consisting of LG 100268, LGD 1069 or 9-cis Retinoic acid.

9. A pharmaceutical composition for the treatment or prevention of hypertriglyceridemia, comprising:

- (a) an RXR agonist;
- (b) a PPAR γ agonist; and
- (c) a pharmaceutically acceptable carrier,

wherein said PPAR γ agonist is a thiazolidinedione compound, and wherein said RXR agonist is selected from the group consisting of LG 100268, LGD 1069 or 9-cis Retinoic acid.

10. A pharmaceutical composition for the treatment of low levels of lipoprotein lipase, comprising:

- (a) an RXR agonist;
- (b) a PPAR γ agonist; and
- (c) a pharmaceutically acceptable carrier,

wherein said PPAR γ agonist is a thiazolidinedione compound, and wherein said RXR agonist is selected from the group consisting of LG 100268, LGD 1069 or 9-cis Retinoic acid.

11. A pharmaceutical composition for the treatment of low levels of HDL cholesterol, comprising:

- (a) an RXR agonist;
- (b) a PPAR γ agonist; and
- (c) a pharmaceutically acceptable carrier,

wherein said PPAR γ agonist is a thiazolidinedione compound, and wherein said RXR agonist is selected from the group consisting of LG 100268, LGD 1069 or 9-cis Retinoic acid.

12. A pharmaceutical composition for increasing adipocyte differentiation, comprising:

- (a) an RXR agonist;
- (b) a PPAR γ agonist; and
- (c) a pharmaceutically acceptable carrier,

wherein said PPAR γ agonist is a thiazolidinedione compound, and wherein said RXR agonist is selected from the group consisting of LG 100268, LGD 1069 or 9-cis Retinoic acid.

13. The composition of any of claims 8-12, wherein said thiazolidinedione compound is selected from the group consisting of BRL 49653, troglitazone, pioglitazone, ciglitazone, WAY-120,744, englitazone, AD 5075 and darglitazone.

14. The composition of claim 13, wherein said composition further comprises insulin, insulin derivative, insulin secretagogue, insulin sensitizer or insulin mimetic.

tagogue, insulin sensitizer or insulin mimetic.

5

10

15

20

25

30


35

40

45

50

55







FIG. 2a.







FIG. 3.








FIG. 4.

FIG. 5.










FIG. 8.





Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 426 048 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:
16.06.2004 Bulletin 2004/25

(43) Date of publication A2:
09.06.2004 Bulletin 2004/24

(21) Application number: 04002911.8

(22) Date of filing: 17.09.1996

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

(30) Priority: 18.09.1995 US 3869 P
06.10.1995 US 4897 P
11.01.1996 US 9884 P
24.05.1996 US 18318 P
10.07.1996 US 21839 P

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
96931613.2 / 0 859 608

(51) Int Cl.7: **A61K 31/192**, A61K 31/4418,
A61K 31/203, A61K 31/426,
A61K 31/4439, A61K 31/455,
A61P 3/06

(71) Applicant: **LIGAND PHARMACEUTICALS, INC.**
San Diego, CA 92121 (US)


(72) Inventors:
• Heyman, Richard A.
Encinitas CA 92024 (US)
• Cesario, Rosemary
San Diego CA 92117 (US)
• Mukherjee, Ranjan
San Diego CA 92127 (US)

(74) Representative: **Viering, Jentschura & Partner**
Steinsdorfstrasse 6
80538 München (DE)

(54) **Treating hypertriglyceridemia with RXR agonists in combination with a PPARgamma agonist**

(57) This invention relates to methods and compositions for the treatment of cardiovascular disease, hypertriglyceridemia, low levels of lipoprotein lipase, low

levels of HDL cholesterol and for increasing adipocyte differentiation using an RXR agonist in combination with a PPARgamma agonist such as a thiazolidinedione compound.





European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 04 00 2911

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	RIGAS J R ET AL: "LIPOPROTEIN ALTERATIONS IN PATIENTS TREATED WITH NOVEL RETINOIDS" PROCEEDINGS OF THE ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH. TORONTO, MAR. 18 - 22, 1995, PHILADELPHIA, AACR, US, vol. MEETING 86, 1 March 1995 (1995-03-01), page 506 XP000615382 ISSN: 0197-016X * the whole document *	1-14	A61K31/192 A61K31/4418 A61K31/203 A61K31/426 A61K31/4439 A61K31/455 A61P3/06
A	US 5 219 888 A (KATOCS JR ANDREW S ET AL) 15 June 1993 (1993-06-15) * the whole document *	1-14	
A	LEHMANN J M ET AL: "AN ANTIDIABETIC THIAZOLIDINEDIONE IS A HIGH AFFINITY LIGAND FOR PEROXISOME PROLIFERATOR-ACTIVATED RECEPTOR GAMMA (PPARGAMMA)" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 270, no. 22, 2 June 1995 (1995-06-02), pages 12953-12956, XP000577082 ISSN: 0021-9258 * the whole document *	1-14	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A61K A61P
-/--			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 13 April 2004	Examiner Hoff, P
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

EPO FORM 1503 03 82 (P/4001)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 04 00 2911

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	BOEHM M F ET AL: "DESIGN AND SYNTHESIS OF POTENT RETINOID X RECEPTOR SELECTIVE LIGANDS THAT INDUCE APOPTOSIS IN LEUKEMIA CELLS" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 38, no. 16, 4 August 1995 (1995-08-04), pages 3146-3155, XP000615461 ISSN: 0022-2623 * the whole document *	1-14	
A	WO 93/21146 A (LIGAND PHARM INC) 28 October 1993 (1993-10-28) * abstract * * page 14 * * page 17 * * page 59, line 15 - page 60, line 21 * * figure 10; example 45 *	1-14	
A	SAFONOVA I ET AL: "FATTY ACIDS AND RETINOID ACT SYNERGISTICALLY ON ADIPOSE CELL DIFFERENTIATION" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 204, no. 2, 28 October 1994 (1994-10-28), pages 498-504, XP000615920 ISSN: 0006-291X * the whole document *	1-14	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 13 April 2004	Examiner Hoff, P
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 04 00 2911

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

13-04-2004

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5219888	A	15-06-1993	NONE	
WO 9321146	A	28-10-1993	AT 195716 T	15-09-2000
			AT 187434 T	15-12-1999
			AT 177733 T	15-04-1999
			AT 256653 T	15-01-2004
			AU 675430 B2	06-02-1997
			AU 4118893 A	18-11-1993
			AU 691477 B2	21-05-1998
			AU 5586894 A	15-08-1994
			AU 5664294 A	28-02-1995
			AU 6225894 A	15-08-1994
			BR 9306284 A	13-01-1998
			BR 9307784 A	14-11-1995
			CA 2133587 A1	28-10-1993
			CA 2153235 A1	21-07-1994
			CA 2153236 A1	09-02-1995
			DE 69324043 D1	22-04-1999
			DE 69324043 T2	01-07-1999
			DE 69327255 D1	13-01-2000
			DE 69327255 T2	30-03-2000
			DE 69329278 D1	28-09-2000
			DE 69329278 T2	09-08-2001
			DE 69333371 D1	29-01-2004
			DK 637297 T3	18-09-2000
			DK 678086 T3	17-04-2000
			DK 678087 T3	27-09-1999
			EP 0637297 A1	08-02-1995
			EP 0678086 A1	25-10-1995
			EP 0678087 A1	25-10-1995
			EP 0983991 A2	08-03-2000
			EP 0983992 A2	08-03-2000
			ES 2149814 T3	16-11-2000
			ES 2139063 T3	01-02-2000
			ES 2129115 T3	01-06-1999
			GR 3032841 T3	31-07-2000
			GR 3030104 T3	30-07-1999
			GR 3034841 T3	28-02-2001
			JP 8511027 T	19-11-1996
			JP 8505852 T	25-06-1996
			JP 8505359 T	11-06-1996
			LU 90803 A9	26-11-2001
			MX 9306627 A1	29-07-1994
			MX 9306629 A1	29-07-1994
			NO 943943 A	21-12-1994
			PT 637297 T	31-01-2001

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 04 00 2911

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

13-04-2004

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9321146 A	PT	678086 T	31-05-2000
	RU	2144913 C1	27-01-2000
	WO	9321146 A1	28-10-1993
	WO	9415901 A1	21-07-1994
	WO	9504036 A1	09-02-1995
	WO	9415902 A1	21-07-1994

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.